

## 14.1 Exercises

- In Example 2 we considered the function  $W = f(T, v)$ , where  $W$  is the wind-chill index,  $T$  is the actual temperature, and  $v$  is the wind speed. A numerical representation is given in Table 1.
  - What is the value of  $f(-15, 40)$ ? What is its meaning?
  - Describe in words the meaning of the question "For what value of  $v$  is  $f(-20, v) = -30$ ?" Then answer the question.
  - Describe in words the meaning of the question "For what value of  $T$  is  $f(T, 20) = -49$ ?" Then answer the question.
  - What is the meaning of the function  $W = f(-5, v)$ ? Describe the behavior of this function.
  - What is the meaning of the function  $W = f(T, 50)$ ? Describe the behavior of this function.
- The *temperature-humidity index*  $I$  (or *humidex*, for short) is the perceived air temperature when the actual temperature is  $T$  and the relative humidity is  $h$ , so we can write  $I = f(T, h)$ . The following table of values of  $I$  is an excerpt from a table compiled by the National Oceanic & Atmospheric Administration.

**TABLE 3** Apparent temperature as a function of temperature and humidity

		Relative humidity (%)					
		20	30	40	50	60	70
Actual temperature (°F)	$T \backslash h$						
	80	77	78	79	81	82	83
	85	82	84	86	88	90	93
	90	87	90	93	96	100	106
	95	93	96	101	107	114	124
	100	99	104	110	120	132	144

- What is the value of  $f(95, 70)$ ? What is its meaning?
  - For what value of  $h$  is  $f(90, h) = 100$ ?
  - For what value of  $T$  is  $f(T, 50) = 88$ ?
  - What are the meanings of the functions  $I = f(80, h)$  and  $I = f(100, h)$ ? Compare the behavior of these two functions of  $h$ .
- A manufacturer has modeled its yearly production function  $P$  (the monetary value of its entire production in millions of dollars) as a Cobb-Douglas function

$$P(L, K) = 1.47L^{0.65}K^{0.35}$$

where  $L$  is the number of labor hours (in thousands) and  $K$  is the invested capital (in millions of dollars). Find  $P(120, 20)$  and interpret it.

- Verify for the Cobb-Douglas production function

$$P(L, K) = 1.01L^{0.75}K^{0.25}$$

discussed in Example 3 that the production will be doubled if both the amount of labor and the amount of capital are doubled. Determine whether this is also true for the general production function

$$P(L, K) = bL^{\alpha}K^{1-\alpha}$$

- A model for the surface area of a human body is given by the function

$$S = f(w, h) = 0.1091w^{0.425}h^{0.725}$$

where  $w$  is the weight (in pounds),  $h$  is the height (in inches), and  $S$  is measured in square feet.

- Find  $f(160, 70)$  and interpret it.
  - What is your own surface area?
- The wind-chill index  $W$  discussed in Example 2 has been modeled by the following function:

$$W(T, v) = 13.12 + 0.6215T - 11.37v^{0.16} + 0.3965Tv^{0.16}$$

Check to see how closely this model agrees with the values in Table 1 for a few values of  $T$  and  $v$ .

- The wave heights  $h$  in the open sea depend on the speed  $v$  of the wind and the length of time  $t$  that the wind has been blowing at that speed. Values of the function  $h = f(v, t)$  are recorded in feet in Table 4.
  - What is the value of  $f(40, 15)$ ? What is its meaning?
  - What is the meaning of the function  $h = f(30, t)$ ? Describe the behavior of this function.
  - What is the meaning of the function  $h = f(v, 30)$ ? Describe the behavior of this function.

**TABLE 4**

		Duration (hours)						
		5	10	15	20	30	40	50
Wind speed (knots)	$v \backslash t$							
	10	2	2	2	2	2	2	2
	15	4	4	5	5	5	5	5
	20	5	7	8	8	9	9	9
	30	9	13	16	17	18	19	19
	40	14	21	25	28	31	33	33
	50	19	29	36	40	45	48	50
	60	24	37	47	54	62	67	69

- A company makes three sizes of cardboard boxes: small, medium, and large. It costs \$2.50 to make a small box, \$4.00

for a medium box, and \$4.50 for a large box. Fixed costs are \$8000.

- (a) Express the cost of making  $x$  small boxes,  $y$  medium boxes, and  $z$  large boxes as a function of three variables:  
 $C = f(x, y, z)$ .

(b) Find  $f(3000, 5000, 4000)$  and interpret it.

(c) What is the domain of  $f$ ?

9. Let  $g(x, y) = \cos(x + 2y)$ .

(a) Evaluate  $g(2, -1)$ .

(b) Find the domain of  $g$ .

(c) Find the range of  $g$ .

10. Let  $F(x, y) = 1 + \sqrt{4 - y^2}$ .

(a) Evaluate  $F(3, 1)$ .

(b) Find and sketch the domain of  $F$ .

(c) Find the range of  $F$ .

11. Let  $f(x, y, z) = \sqrt{x} + \sqrt{y} + \sqrt{z} + \ln(4 - x^2 - y^2 - z^2)$ .

(a) Evaluate  $f(1, 1, 1)$ .

(b) Find and describe the domain of  $f$ .

12. Let  $g(x, y, z) = x^2y^2z\sqrt{10 - x - y - z}$ .

(a) Evaluate  $g(1, 2, 3)$ .

(b) Find and describe the domain of  $g$ .

13–22 Find and sketch the domain of the function.

13.  $f(x, y) = \sqrt{2x - y}$

14.  $f(x, y) = \sqrt{xy}$

15.  $f(x, y) = \ln(9 - x^2 - 9y^2)$

16.  $f(x, y) = \sqrt{x^2 - y^2}$

17.  $f(x, y) = \sqrt{1 - x^2} - \sqrt{1 - y^2}$

18.  $f(x, y) = \sqrt{y} + \sqrt{25 - x^2 - y^2}$

19.  $f(x, y) = \frac{\sqrt{y - x^2}}{1 - x^2}$

20.  $f(x, y) = \arcsin(x^2 + y^2 - 2)$

21.  $f(x, y, z) = \sqrt{1 - x^2 - y^2 - z^2}$

22.  $f(x, y, z) = \ln(16 - 4x^2 - 4y^2 - z^2)$

23–31 Sketch the graph of the function.

23.  $f(x, y) = 1 + y$

24.  $f(x, y) = 2 - x$

25.  $f(x, y) = 10 - 4x - 5y$

26.  $f(x, y) = e^{-y}$

27.  $f(x, y) = y^2 + 1$

28.  $f(x, y) = 1 + 2x^2 + 2y^2$

29.  $f(x, y) = 9 - x^2 - 9y^2$

30.  $f(x, y) = \sqrt{4x^2 + y^2}$

31.  $f(x, y) = \sqrt{4 - 4x^2 - y^2}$

32. Match the function with its graph (labeled I–VI). Give reasons for your choices.

(a)  $f(x, y) = |x| + |y|$


(b)  $f(x, y) = |xy|$

(c)  $f(x, y) = \frac{1}{1 + x^2 + y^2}$


(d)  $f(x, y) = (x^2 - y^2)^2$

(e)  $f(x, y) = (x - y)^2$

(f)  $f(x, y) = \sin(|x| + |y|)$




33. A contour map for a function  $f$  is shown. Use it to estimate the values of  $f(-3, 3)$  and  $f(3, -2)$ . What can you say about the shape of the graph?




34. Shown is a contour map of atmospheric pressure in North America on August 12, 2008. On the level curves (called isobars) the pressure is indicated in millibars (mb).

- (a) Estimate the pressure at  $C$  (Chicago),  $N$  (Nashville),  $S$  (San Francisco), and  $V$  (Vancouver).


(b) At which of these locations were the winds strongest?



35. Level curves (isothermals) are shown for the water temperature (in °C) in Long Lake (Minnesota) in 1998 as a function of depth and time of year. Estimate the temperature in the lake on June 9 (day 160) at a depth of 10 m and on June 29 (day 180) at a depth of 5 m.




36. Two contour maps are shown. One is for a function  $f$  whose graph is a cone. The other is for a function  $g$  whose graph is a paraboloid. Which is which, and why?




37. Locate the points  $A$  and  $B$  on the map of Lonesome Mountain (Figure 12). How would you describe the terrain near  $A$ ? Near  $B$ ?

38. Make a rough sketch of a contour map for the function whose graph is shown.



- 39–42 A contour map of a function is shown. Use it to make a rough sketch of the graph of  $f$ .



- 43–50 Draw a contour map of the function showing several level curves.

43.  $f(x, y) = (y - 2x)^2$       44.  $f(x, y) = x^3 - y$   
 45.  $f(x, y) = \sqrt{x} + y$       46.  $f(x, y) = \ln(x^2 + 4y^2)$   
 47.  $f(x, y) = ye^x$       48.  $f(x, y) = y \sec x$   
 49.  $f(x, y) = \sqrt{y^2 - x^2}$       50.  $f(x, y) = y/(x^2 + y^2)$


- 51–52 Sketch both a contour map and a graph of the function and compare them.

51.  $f(x, y) = x^2 + 9y^2$       52.  $f(x, y) = \sqrt{36 - 9x^2 - 4y^2}$

53. A thin metal plate, located in the  $xy$ -plane, has temperature  $T(x, y)$  at the point  $(x, y)$ . The level curves of  $T$  are called *isothermals* because at all points on such a curve the temperature is the same. Sketch some isothermals if the temperature function is given by

$$T(x, y) = \frac{100}{1 + x^2 + 2y^2}$$

54. If  $V(x, y)$  is the electric potential at a point  $(x, y)$  in the  $xy$ -plane, then the level curves of  $V$  are called *equipotential curves* because at all points on such a curve the electric potential is the same. Sketch some equipotential curves if  $V(x, y) = c/\sqrt{r^2 - x^2 - y^2}$ , where  $c$  is a positive constant.

 **55–58** Use a computer to graph the function using various domains and viewpoints. Get a printout of one that, in your opinion, gives a good view. If your software also produces level curves, then plot some contour lines of the same function and compare with the graph.

**55.**  $f(x, y) = xy^2 - x^3$  (monkey saddle)

**56.**  $f(x, y) = xy^3 - yx^3$  (dog saddle)

**57.**  $f(x, y) = e^{-(x^2+y^2)/3}(\sin(x^2) + \cos(y^2))$

**58.**  $f(x, y) = \cos x \cos y$

**59–64** Match the function (a) with its graph (labeled A–F below) and (b) with its contour map (labeled I–VI). Give reasons for your choices.

**59.**  $z = \sin(xy)$













**60.**  $z = e^x \cos y$

**61.**  $z = \sin(x - y)$

**62.**  $z = \sin x - \sin y$

**63.**  $z = (1 - x^2)(1 - y^2)$

**64.**  $z = \frac{x - y}{1 + x^2 + y^2}$



65–68 Describe the level surfaces of the function.

65.  $f(x, y, z) = x + 3y + 5z$

66.  $f(x, y, z) = x^2 + 3y^2 + 5z^2$

67.  $f(x, y, z) = y^2 + z^2$

68.  $f(x, y, z) = x^2 - y^2 - z^2$

69–70 Describe how the graph of  $g$  is obtained from the graph of  $f$ .

69. (a)  $g(x, y) = f(x, y) + 2$

(b)  $g(x, y) = 2f(x, y)$

(c)  $g(x, y) = -f(x, y)$

(d)  $g(x, y) = 2 - f(x, y)$

70. (a)  $g(x, y) = f(x - 2, y)$

(b)  $g(x, y) = f(x, y + 2)$

(c)  $g(x, y) = f(x + 3, y - 4)$

71–72 Use a computer to graph the function using various domains and viewpoints. Get a printout that gives a good view of the “peaks and valleys.” Would you say the function has a maximum value? Can you identify any points on the graph that you might consider to be “local maximum points”? What about “local minimum points”?

71.  $f(x, y) = 3x - x^4 - 4y^2 - 10xy$

72.  $f(x, y) = xy e^{-x^2-y^2}$

73–74 Use a computer to graph the function using various domains and viewpoints. Comment on the limiting behavior of the function. What happens as both  $x$  and  $y$  become large? What happens as  $(x, y)$  approaches the origin?

73.  $f(x, y) = \frac{x + y}{x^2 + y^2}$

74.  $f(x, y) = \frac{xy}{x^2 + y^2}$

75. Use a computer to investigate the family of functions  $f(x, y) = e^{cx^2+xy}$ . How does the shape of the graph depend on  $c$ ?

76. Use a computer to investigate the family of surfaces

$$z = (ax^2 + by^2)e^{-x^2-y^2}$$

How does the shape of the graph depend on the numbers  $a$  and  $b$ ?

77. Use a computer to investigate the family of surfaces  $z = x^2 + y^2 + cxy$ . In particular, you should determine the transitional values of  $c$  for which the surface changes from one type of quadric surface to another.

78. Graph the functions

$$f(x, y) = \sqrt{x^2 + y^2}$$

$$f(x, y) = e^{\sqrt{x^2+y^2}}$$

$$f(x, y) = \ln\sqrt{x^2 + y^2}$$

$$f(x, y) = \sin(\sqrt{x^2 + y^2})$$

and 
$$f(x, y) = \frac{1}{\sqrt{x^2 + y^2}}$$

In general, if  $g$  is a function of one variable, how is the graph of

$$f(x, y) = g(\sqrt{x^2 + y^2})$$

obtained from the graph of  $g$ ?

79. (a) Show that, by taking logarithms, the general Cobb-Douglas function  $P = bL^\alpha K^{1-\alpha}$  can be expressed as

$$\ln \frac{P}{K} = \ln b + \alpha \ln \frac{L}{K}$$

- (b) If we let  $x = \ln(L/K)$  and  $y = \ln(P/K)$ , the equation in part (a) becomes the linear equation  $y = \alpha x + \ln b$ . Use Table 2 (in Example 3) to make a table of values of  $\ln(L/K)$  and  $\ln(P/K)$  for the years 1899–1922. Then use a graphing calculator or computer to find the least squares regression line through the points  $(\ln(L/K), \ln(P/K))$ .
- (c) Deduce that the Cobb-Douglas production function is  $P = 1.01L^{0.75}K^{0.25}$ .