

## 6.5 Exercises


---


1. A population of protozoa develops with a constant relative growth rate of 0.7944 per member per day. On day zero the population consists of two members. Find the population size after six days.
2. A common inhabitant of human intestines is the bacterium *Escherichia coli*. A cell of this bacterium in a nutrient-broth medium divides into two cells every 20 minutes. The initial population of a culture is 60 cells.
  - (a) Find the relative growth rate.
  - (b) Find an expression for the number of cells after  $t$  hours.
  - (c) Find the number of cells after 8 hours.
  - (d) Find the rate of growth after 8 hours.
  - (e) When will the population reach 20,000 cells?
3. A bacteria culture initially contains 100 cells and grows at a rate proportional to its size. After an hour the population has increased to 420.
  - (a) Find an expression for the number of bacteria after  $t$  hours.
  - (b) Find the number of bacteria after 3 hours.
  - (c) Find the rate of growth after 3 hours.
  - (d) When will the population reach 10,000?



4. A bacteria culture grows with constant relative growth rate. The bacteria count was 400 after 2 hours and 25,600 after 6 hours.
- What is the relative growth rate? Express your answer as a percentage.
  - What was the initial size of the culture?
  - Find an expression for the number of bacteria after  $t$  hours.
  - Find the number of cells after 4.5 hours.
  - Find the rate of growth after 4.5 hours.
  - When will the population reach 50,000?
5. The table gives estimates of the world population, in millions, from 1750 to 2000.
- Use the exponential model and the population figures for 1750 and 1800 to predict the world population in 1900 and 1950. Compare with the actual figures.
  - Use the exponential model and the population figures for 1850 and 1900 to predict the world population in 1950. Compare with the actual population.
  - Use the exponential model and the population figures for 1900 and 1950 to predict the world population in 2000. Compare with the actual population and try to explain the discrepancy.

Year	Population	Year	Population
1750	790	1900	1650
1800	980	1950	2560
1850	1260	2000	6080

6. The table gives the population of India, in millions, for the second half of the 20th century.
- | Year | Population |
|------|------------|
| 1951 | 361        |
| 1961 | 439        |
| 1971 | 548        |
| 1981 | 683        |
| 1991 | 846        |
| 2001 | 1029       |
- Use the exponential model and the census figures for 1951 and 1961 to predict the population in 2001. Compare with the actual figure.
  - Use the exponential model and the census figures for 1961 and 1981 to predict the population in 2001. Compare with the actual population. Then use this model to predict the population in the years 2010 and 2020.
- 
  - Graph both of the exponential functions in parts (a) and (b) together with a plot of the actual population. Are these models reasonable ones?
7. Experiments show that if the chemical reaction



takes place at  $45^\circ\text{C}$ , the rate of reaction of dinitrogen pent-

oxide is proportional to its concentration as follows:

$$-\frac{d[\text{N}_2\text{O}_5]}{dt} = 0.0005[\text{N}_2\text{O}_5]$$

(See Example 4 in Section 2.7.)

- Find an expression for the concentration  $[\text{N}_2\text{O}_5]$  after  $t$  seconds if the initial concentration is  $C$ .
  - How long will the reaction take to reduce the concentration of  $\text{N}_2\text{O}_5$  to 90% of its original value?
8. Strontium-90 has a half-life of 28 days.
- A sample has a mass of 50 mg initially. Find a formula for the mass remaining after  $t$  days.
  - Find the mass remaining after 40 days.
  - How long does it take the sample to decay to a mass of 2 mg?
  - Sketch the graph of the mass function.
9. The half-life of cesium-137 is 30 years. Suppose we have a 100-mg sample.
- Find the mass that remains after  $t$  years.
  - How much of the sample remains after 100 years?
  - After how long will only 1 mg remain?
10. A sample of tritium-3 decayed to 94.5% of its original amount after a year.
- What is the half-life of tritium-3?
  - How long would it take the sample to decay to 20% of its original amount?
11. Scientists can determine the age of ancient objects by the method of *radiocarbon dating*. The bombardment of the upper atmosphere by cosmic rays converts nitrogen to a radioactive isotope of carbon,  $^{14}\text{C}$ , with a half-life of about 5730 years. Vegetation absorbs carbon dioxide through the atmosphere and animal life assimilates  $^{14}\text{C}$  through food chains. When a plant or animal dies, it stops replacing its carbon and the amount of  $^{14}\text{C}$  begins to decrease through radioactive decay. Therefore the level of radioactivity must also decay exponentially.
- A parchment fragment was discovered that had about 74% as much  $^{14}\text{C}$  radioactivity as does plant material on the earth today. Estimate the age of the parchment.
12. A curve passes through the point  $(0, 5)$  and has the property that the slope of the curve at every point  $P$  is twice the  $y$ -coordinate of  $P$ . What is the equation of the curve?
13. A roast turkey is taken from an oven when its temperature has reached  $185^\circ\text{F}$  and is placed on a table in a room where the temperature is  $75^\circ\text{F}$ .
- If the temperature of the turkey is  $150^\circ\text{F}$  after half an hour, what is the temperature after 45 minutes?
  - When will the turkey have cooled to  $100^\circ\text{F}$ ?
14. In a murder investigation, the temperature of the corpse was  $32.5^\circ\text{C}$  at 1:30 PM and  $30.3^\circ\text{C}$  an hour later. Normal body temperature is  $37.0^\circ\text{C}$  and the temperature of the surroundings was  $20.0^\circ\text{C}$ . When did the murder take place?

15. When a cold drink is taken from a refrigerator, its temperature is  $5^{\circ}\text{C}$ . After 25 minutes in a  $20^{\circ}\text{C}$  room its temperature has increased to  $10^{\circ}\text{C}$ .

- (a) What is the temperature of the drink after 50 minutes?  
 (b) When will its temperature be  $15^{\circ}\text{C}$ ?

16. (a) A cup of coffee has temperature  $95^{\circ}\text{C}$  and takes 30 minutes to cool to  $61^{\circ}\text{C}$  in a room with temperature  $20^{\circ}\text{C}$ . Show that the temperature of the coffee after  $t$  minutes is

$$T(t) = 20 + 75e^{-kt}$$

where  $k \approx 0.02$ .

- (b) What is the average temperature of the coffee during the first half hour?

17. The rate of change of atmospheric pressure  $P$  with respect to altitude  $h$  is proportional to  $P$ , provided that the temperature is constant. At  $15^{\circ}\text{C}$  the pressure is 101.3 kPa at sea level and 87.14 kPa at  $h = 1000$  m.

- (a) What is the pressure at an altitude of 3000 m?  
 (b) What is the pressure at the top of Mount McKinley, at an altitude of 6187 m?

18. (a) If \$1000 is borrowed at 8% interest, find the amounts due at the end of 3 years if the interest is compounded (i) annually, (ii) quarterly, (iii) monthly, (iv) weekly, (v) daily, (vi) hourly, and (vii) continuously.



(b) Suppose \$1000 is borrowed and the interest is compounded continuously. If  $A(t)$  is the amount due after  $t$  years, where  $0 \leq t \leq 3$ , graph  $A(t)$  for each of the interest rates 6%, 8%, and 10% on a common screen.

19. (a) If \$3000 is invested at 5% interest, find the value of the investment at the end of 5 years if the interest is compounded (i) annually, (ii) semiannually, (iii) monthly, (iv) weekly, (v) daily, and (vi) continuously.

(b) If  $A(t)$  is the amount of the investment at time  $t$  for the case of continuous compounding, write a differential equation and an initial condition satisfied by  $A(t)$ .

20. (a) How long will it take an investment to double in value if the interest rate is 6% compounded continuously?

(b) What is the equivalent annual interest rate?