

## 1.4 Exercises

1. A tank holds 1000 gallons of water, which drains from the bottom of the tank in half an hour. The values in the table show the volume  $V$  of water remaining in the tank (in gallons) after  $t$  minutes.

$t$ (min)	5	10	15	20	25	30
$V$ (gal)	694	444	250	111	28	0

- (a) If  $P$  is the point  $(15, 250)$  on the graph of  $V$ , find the slopes of the secant lines  $PQ$  when  $Q$  is the point on the graph with  $t = 5, 10, 20, 25$ , and  $30$ .  
 (b) Estimate the slope of the tangent line at  $P$  by averaging the slopes of two secant lines.  
 (c) Use a graph of the function to estimate the slope of the tangent line at  $P$ . (This slope represents the rate at which the water is flowing from the tank after 15 minutes.)
2. A cardiac monitor is used to measure the heart rate of a patient after surgery. It compiles the number of heartbeats after  $t$  minutes. When the data in the table are graphed, the slope of the tangent line represents the heart rate in beats per minute.

$t$ (min)	36	38	40	42	44
Heartbeats	2530	2661	2806	2948	3080

The monitor estimates this value by calculating the slope of a secant line. Use the data to estimate the patient's heart rate after 42 minutes using the secant line between the points with the given values of  $t$ .

- (a)  $t = 36$  and  $t = 42$       (b)  $t = 38$  and  $t = 42$   
 (c)  $t = 40$  and  $t = 42$       (d)  $t = 42$  and  $t = 44$

What are your conclusions?

3. The point  $P(2, -1)$  lies on the curve  $y = 1/(1 - x)$ .  
 (a) If  $Q$  is the point  $(x, 1/(1 - x))$ , use your calculator to find the slope of the secant line  $PQ$  (correct to six decimal places) for the following values of  $x$ :  
 (i) 1.5    (ii) 1.9    (iii) 1.99    (iv) 1.999  
 (v) 2.5    (vi) 2.1    (vii) 2.01    (viii) 2.001  
 (b) Using the results of part (a), guess the value of the slope of the tangent line to the curve at  $P(2, -1)$ .  
 (c) Using the slope from part (b), find an equation of the tangent line to the curve at  $P(2, -1)$ .
4. The point  $P(0.5, 0)$  lies on the curve  $y = \cos \pi x$ .  
 (a) If  $Q$  is the point  $(x, \cos \pi x)$ , use your calculator to find the slope of the secant line  $PQ$  (correct to six decimal places) for the following values of  $x$ :  
 (i) 0    (ii) 0.4    (iii) 0.49    (iv) 0.499  
 (v) 1    (vi) 0.6    (vii) 0.51    (viii) 0.501  
 (b) Using the results of part (a), guess the value of the slope of the tangent line to the curve at  $P(0.5, 0)$ .

- (c) Using the slope from part (b), find an equation of the tangent line to the curve at  $P(0.5, 0)$ .  
 (d) Sketch the curve, two of the secant lines, and the tangent line.

5. If a ball is thrown into the air with a velocity of 40 ft/s, its height in feet  $t$  seconds later is given by  $y = 40t - 16t^2$ .  
 (a) Find the average velocity for the time period beginning when  $t = 2$  and lasting  
 (i) 0.5 second    (ii) 0.1 second  
 (iii) 0.05 second    (iv) 0.01 second  
 (b) Estimate the instantaneous velocity when  $t = 2$ .
6. If a rock is thrown upward on the planet Mars with a velocity of 10 m/s, its height in meters  $t$  seconds later is given by  $y = 10t - 1.86t^2$ .  
 (a) Find the average velocity over the given time intervals:  
 (i)  $[1, 2]$     (ii)  $[1, 1.5]$     (iii)  $[1, 1.1]$   
 (iv)  $[1, 1.01]$     (v)  $[1, 1.001]$   
 (b) Estimate the instantaneous velocity when  $t = 1$ .

7. The table shows the position of a cyclist.

$t$ (seconds)	0	1	2	3	4	5
$s$ (meters)	0	1.4	5.1	10.7	17.7	25.8

- (a) Find the average velocity for each time period:  
 (i)  $[1, 3]$     (ii)  $[2, 3]$     (iii)  $[3, 5]$     (iv)  $[3, 4]$   
 (b) Use the graph of  $s$  as a function of  $t$  to estimate the instantaneous velocity when  $t = 3$ .
8. The displacement (in centimeters) of a particle moving back and forth along a straight line is given by the equation of motion  $s = 2 \sin \pi t + 3 \cos \pi t$ , where  $t$  is measured in seconds.  
 (a) Find the average velocity during each time period:  
 (i)  $[1, 2]$     (ii)  $[1, 1.1]$   
 (iii)  $[1, 1.01]$     (iv)  $[1, 1.001]$   
 (b) Estimate the instantaneous velocity of the particle when  $t = 1$ .
9. The point  $P(1, 0)$  lies on the curve  $y = \sin(10\pi/x)$ .  
 (a) If  $Q$  is the point  $(x, \sin(10\pi/x))$ , find the slope of the secant line  $PQ$  (correct to four decimal places) for  $x = 2, 1.5, 1.4, 1.3, 1.2, 1.1, 0.5, 0.6, 0.7, 0.8$ , and  $0.9$ . Do the slopes appear to be approaching a limit?  
 (b) Use a graph of the curve to explain why the slopes of the secant lines in part (a) are not close to the slope of the tangent line at  $P$ .  
 (c) By choosing appropriate secant lines, estimate the slope of the tangent line at  $P$ .

